arXiv:1605.02728v1 [math.CA] 9 May 2016

Integrals and Series Representations of q-Polynomials and Functions: Part II Schur Polynomials and the Rogers-Ramanujan Identities

Mourad E. H. Ismail * Ruiming Zhang †

May 11, 2016

Abstract

We give several expansion and identities involving the Ramanujan function A_q and the Stieltjes–Wigert polynomials. Special values of our idenitities give m-versions of some of the items on the Slater list of Rogers-Ramanujan type identities. We also study some bilateral extensions of certain transformations in the theory of basic hypergeometric functions.

Mathematics Subject Classification MSC 2010: Primary 11P84, 33D45 Secondary 05A17.

Keywords and phrases: Rogers-Ramanujan identities, m-versions, the Ramanujan function, Stieltjes–Wigert polynomials, bilateral q-series.

Filename: IsmZhaRRII1

1 Introduction

In part I of this series of papers we derived several integral representations for many q-functions and polynomials including the the q-exponential functions e_q, E_q, and our $\mathcal{E}_q$, [20]. We also derived several series identities and transformations. The present work is part II where we continue our studies and establish quite a few series identities and transformation formulas. Some of our formulas give new identities for the Schur polynomials introduced by I. Schur in [28]. We also treat a generalization of the Schur polynomials we introduced in [22].

We recall the Rogers–Ramanujan identities are

$$\sum_{n=0}^{\infty}\frac{q^{n^2}}{(q;q)_n}=\frac{1}{(q,q^4;q^5)_\infty},\qquad \sum_{n=0}^{\infty}\frac{q^{n^2+n}}{(q;q)_n}=\frac{1}{(q^2,q^3;q^5)_\infty}, \tag{1.1}$$

where the notation for the q-shifted factorials is the standard notation followed in [12], [5], or [15]. References for the Rogers-Ramanujan identities, their origins and many of their applications are in [2], [3], and [5]. Garrett, Ismail, and Stanton [11] generalized the Rogers–Ramanujan to

$$\sum_{n=0}^{\infty}\frac{q^{n^2+mn}}{(q;q)_n}=\frac{(-1)^m q^{-\binom{m}{2}}a_m(q)}{(q,q^4;q^5)_\infty}-\frac{(-1)^m q^{-\binom{m}{2}}b_m(q)}{(q^2,q^3;q^5)_\infty}, \tag{1.2}$$

*Research partially supported by the DSFP of King Saud University and by the National Plan for Science, Technology and innovation (MAARIFAH), King Abdelaziz City for Science and Technology, Kingdom of Saudi Arabia, Award number 14-MAT623-02.

†Corresponding author, research partially supported by National Science Foundation of China, grant No. 11371294.

where $a_m(q)$ and $b_m(q)$ are defined by

$$a_m(q) = \sum_{j\geq 0} q^{j^2+j} \begin{bmatrix} m-j-2 \\ j \end{bmatrix}_q, \qquad b_m(q) = \sum_{j\geq 0} q^{j^2} \begin{bmatrix} m-j-1 \\ j \end{bmatrix}_q. \tag{1.3}$$

The Garrett–Ismail–Stanton result became known as the m-version of the Rogers-Ramanujan identities.

The polynomials $a_m(q)$ and $b_m(q)$ were considered by Schur in conjunction with his proof of the Rogers–Ramanujan identities, see [2] and [11] for details. We shall refer to $a_m(q)$ and $b_m(q)$ as the Schur polynomials. The closed form expressions for a_m and b_m in (1.3) were given by Andrews in [4], where he also gave a polynomial generalization of the Rogers–Ramanujan identities. We must note that $a_{m+1}(q)$ and $b_{m+1}(q)$ are the partial numberators and denominators of the Ramanujan continued fraction,

In this work we will use the following confluent limit of the q-Gauus sum

$${}_1\phi_1\left(a;c;q,c/a\right) = \frac{(c/a;q)_\infty}{(c;q)_\infty} \tag{1.4}$$

We will also use the integral representations,

$$\begin{aligned} &(bq^\alpha;q)_\infty\, q^{\alpha^2/2}{}_1\phi_1\left(a;bq^\alpha;q,zq^{\alpha+1/2}\right) \\ &= \frac{1}{\sqrt{\pi\log q^{-2}}}\int_{-\infty}^{\infty} \frac{\left(-aze^{ix};q\right)_\infty \exp\left(\frac{x^2}{\log q^2}+i\alpha x\right)dx}{\left(-bq^{-1/2}e^{ix},-ze^{ix};q\right)_\infty}, \end{aligned} \tag{1.5}$$

$$q^{\alpha^2/2}A_q\left(q^\alpha z\right) = \frac{1}{\sqrt{2\pi}}\int_{-\infty}^{\infty} \frac{\left(zq^{1/2}e^{ix};q\right)_\infty \exp\left(\frac{x^2}{\log q^2}+i\alpha x\right)}{\sqrt{\log q^{-1}}}dx, \tag{1.6}$$

$$q^{\alpha^2/2}\left(-zq^{\alpha+1/2};q\right)_\infty = \int_{-\infty}^{\infty} \frac{\exp\left(x^2/\log q^2+i\alpha x\right)dx}{\left(ze^{ix};q\right)_\infty\sqrt{\pi\log q^{-2}}}, \tag{1.7}$$

and

$$q^{\alpha^2/2}S_n\left(xq^{\alpha-1/2};q\right) = \int_{-\infty}^{\infty} \frac{\left(xe^{iy};q\right)_n}{(q;q)_n} \frac{\exp\left(y^2/\log q^2+i\alpha y\right)dy}{\sqrt{\pi\log q^{-2}}}. \tag{1.8}$$

which we proved in our forthcoming paper [23].

In Section 2 we prove the following theorem and discuss some of its implications.

Theorem 1.1. *We have the indentites*

$$(-q;q)_\infty = \frac{1}{(q,q^4;q^5)_\infty}\sum_{n=0}^{\infty}\frac{q^{2n}a_{2n}(q)}{(q^2;q^2)_n} - \frac{1}{(q^2,q^3;q^5)_\infty}\sum_{n=0}^{\infty}\frac{q^{2n}b_{2n}(q)}{(q^2;q^2)_n}. \tag{1.9}$$

$$(-q^2;q)_\infty = \frac{1}{(q^2,q^3;q^5)_\infty}\sum_{n=0}^{\infty}\frac{q^{2n}b_{2n+1}(q)}{(q^2;q^2)_n} - \frac{1}{(q,q^4;q^5)_\infty}\sum_{n=0}^{\infty}\frac{q^{2n}a_{2n+1}(q)}{(q^2;q^2)_n}. \tag{1.10}$$

This theorem will follow from Theorem 1.2 which we now state.

Theorem 1.2. *The following identities hold for all $a, z \in \mathbb{C}$ but $z \neq q^n, n = 0, -1, -2, \cdots$,*

$$(z;q)_\infty \, {}_1\phi_1 \left(a; z; q, -z\right) = \sum_{n=0}^{\infty} \frac{q^{2n^2-n} z^{2n}}{\left(q^2;q^2\right)_n} A_q\left(q^{2n-1} a z\right), \tag{1.11}$$

$$(b;q)_\infty \, {}_1\phi_1 \left(a; b; q, -b\right) = \sum_{n=0}^{\infty} \frac{q^{n^2-n} \left(-ab\right)^n}{\left(q;q\right)_n} A_{q^2}\left(-q^{2n-1} b^2\right), \tag{1.12}$$

$$\sum_{n=0}^{\infty} \frac{q^{2n^2-n} b^{2n}}{\left(q^2;q^2\right)_n} A_q\left(q^{2n-1} a b\right) = \sum_{n=0}^{\infty} \frac{q^{n^2-n} \left(-ab\right)^n}{\left(q;q\right)_n} A_{q^2}\left(-q^{2n-1} b^2\right). \tag{1.13}$$

Formula (1.11) is (7.32) in our paper [22] while (1.12) is the result of writing $A_q(q^{2n-1}az)$ as a sum then interchange the two sums, which is then stated formally as (1.13).

Corollary 1.3. *We have*

$$(-z;q)_\infty = \sum_{n=0}^{\infty} \frac{q^{2n^2-n} z^{2n}}{\left(q^2;q^2\right)_n} A_q\left(-q^{2n-1} z\right), \tag{1.14}$$

This is just the case $a = 1$ of (1.11). Recall the definition of the Jackson modified q-Bessel functions, using the notation from, [13],

$$I_\nu^{(1)}(z;q) = \frac{\left(q^{\nu+1};q\right)_\infty}{\left(q;q\right)_\infty} \sum_{n=0}^{\infty} \frac{(z/2)^{\nu+2n}}{\left(q, q^{\nu+1};q\right)_n}, \quad |z| < 2, \tag{1.15}$$

$$I_\nu^{(2)}(z;q) = \frac{\left(q^{\nu+1};q\right)_\infty}{\left(q;q\right)_\infty} \sum_{n=0}^{\infty} \frac{q^{n(n+\nu)}}{\left(q, q^{\nu+1};q\right)_n} (z/2)^{\nu+2n}, \tag{1.16}$$

$$I_\nu^{(3)}(z;q) = \frac{\left(q^{\nu+1};q\right)_\infty}{\left(q;q\right)_\infty} \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}}}{\left(q, q^{\nu+1};q\right)_n} (z/2)^{\nu+2n}. \tag{1.17}$$

In our work [23] we proved Theorem 1.4 below.

Theorem 1.4. *We have the inverse pair*

$$J_\nu^{(2)}\left(2z;q\right) = \frac{z^\nu}{\left(q;q\right)_\infty} \sum_{k=0}^{\infty} \frac{\left(-q^\nu\right)^k}{\left(q;q\right)_k} q^{\binom{k+1}{2}} A_q\left(q^{\nu+k} z^2\right), \tag{1.18}$$

$$\frac{z^\nu A_q\left(q^\nu z^2\right)}{\left(q;q\right)_\infty} = \sum_{k=0}^{\infty} \frac{q^{k^2}}{\left(q;q\right)_k} \left(\frac{q^\nu}{z}\right)^k J_{k+\nu}^{(2)}\left(2z;q\right). \tag{1.19}$$

In the same work we also proved that

$$\frac{A_q\left(z\right)}{\left(zq^2;q^2\right)_\infty} = \sum_{k=0}^{\infty} \frac{\left(-z\right)^k q^{k^2}}{\left(q^2, zq^2;q^2\right)_k}. \tag{1.20}$$

$$S_n\left(x;q\right) = \frac{1}{\left(q;q\right)_n} \sum_{k=0}^{\infty} \frac{\left(xq^n\right)^k}{\left(q;q\right)_k} q^{\binom{k+1}{2}} A_q\left(q^k x\right), \tag{1.21}$$

$$A_q\left(ab\right) = \sum_{k=0}^{\infty} \frac{\left(b;q\right)_k}{\left(q;q\right)_k} q^{\binom{k+1}{2}} a^k A_q\left(aq^k\right), \tag{1.22}$$

see equations (7.6), (7.8), and (7.2), respectively, in [23]. The Stieltjes–Wigert polynomials S_n in (1.21) are defined by

$$S_n(x) = \sum_{k=0}^{n} \frac{q^{k^2}}{(q;q)_k (q;q)_{n-k}} (-z)^k. \tag{1.23}$$

In Section 2 we also discuss some consequences of the results stated so far. This leads, among other things, to new generating functions for the Schur polynomials. Section 3 contains a master identity for bilateral q-series and some noteworthy special and limiting cases of it. Section 4 has an extensive list of new identities involving the Ramanujan function and special cases of them lead to m-versions of formulas on the Slater list [29]. It also contains several new results on the Stieltjes–Wigert polynomials.

2 Theorems 1.1-1.4

It is obvious that (1.2) is nothing but

$$A_q(-q^m)=\sum_{n=0}^{\infty}\frac{q^{n^2+mn}}{(q;q)_n}=\frac{(-1)^m q^{-\binom{m}{2}}a_m(q)}{(q,q^4;q^5)_\infty}-\frac{(-1)^m q^{-\binom{m}{2}}b_m(q)}{(q^2,q^3;q^5)_\infty}. \tag{2.1}$$

This formula will be used repeatedly in this work.

Corollary 1.3 is the case $a=1$ of Theorem 1.2.

Proof of Theorem 1.1. We take $z=q$ in Cor 1.3 and apply (1.2) to obtain

$$\begin{aligned}(-q;q)_\infty &= \sum_{n=0}^{\infty}\frac{q^{2n^2+n}}{(q^2;q^2)_n}A_q\left(-q^{2n}\right)\\ &= \sum_{n=0}^{\infty}\frac{q^{2n^2+n}}{(q^2;q^2)_n}\left(\frac{q^{-\binom{2n}{2}}a_{2n}(q)}{(q,q^4;q^5)_\infty}-\frac{q^{-\binom{2n}{2}}b_{2n}(q)}{(q^2,q^3;q^5)_\infty}\right),\end{aligned}$$

which simplifies to (1.9). Similarly $z=q^2$ in (1.14) leads to (1.10). □

Let $a=0$ in (1.11) to get

$$(z;q)_\infty\sum_{n=0}^{\infty}\frac{q^{\binom{n}{2}}z^n}{(q,z;q)_n}=\sum_{n=0}^{\infty}\frac{q^{2n^2-n}z^{2n}}{(q^2;q^2)_n}.$$

Obviously the above equation is

$$A_{q^2}(-z^2/q)=(z;q)_\infty\sum_{n=0}^{\infty}\frac{q^{\binom{n}{2}}z^n}{(q,z;q)_n}, \tag{2.2}$$

which relates the Ramanujan function to a modified q-Bessel function $I_\nu^{(3)}$. The special case $z=q^{m+1/2}$ is of interest and gives

$$\begin{aligned}&\left(q^{m+1/2};q\right)_\infty\sum_{n=0}^{\infty}\frac{q^{mn+n^2/2}}{(q,q^{m+1/2};q)_n}\\ &\quad=(-1)^m q^{2\binom{m}{2}}\left[\frac{a_m(q^2)}{(q^2,q^8;q^{10})_\infty}-\frac{b_m(q^2)}{(q^4,q^6;q^{10})_\infty}\right].\end{aligned} \tag{2.3}$$

It is clear that the left-hand side is a q-analogue of the fact that $I_{m+1/2}$ is a linear combination of $I_{\pm 1/2}$ with coefficients related to the Lommel polynomials. The cases $m=0,1$ are

$$\sum_{n=0}^{\infty}\frac{q^{n^2}}{(q;q)_{2n}} = \frac{1}{(q;q^2)_\infty\,(q^4,q^{16};q^{20})_\infty}, \tag{2.4}$$

$$\sum_{n=0}^{\infty}\frac{q^{n(n+2)}}{(q;q)_{2n}} = \frac{1}{(q;q^3)_\infty\,(q^8,q^{12};q^{20})_\infty}, \tag{2.5}$$

The identity (2.4) is (79) on Lucy Slater's list, [29], but (2.5) is not on her list.

For $\ell \in \mathbb{N}_0$, we let $a = -q^{\ell+1}/z$ in (1.11) and establish

$$(z;q)_\infty \, {}_1\phi_1\left(-q^{\ell+1}/z; z; q, -z\right) = \sum_{n=0}^{\infty} \frac{q^{2n^2-n} z^{2n}}{(q^2;q^2)_n} A_q\left(-q^{2n+\ell}\right)$$

$$= \frac{(-1)^\ell q^{-\binom{\ell}{2}}}{(q, q^4; q^5)_\infty} \sum_{n=0}^{\infty} \frac{a_{2n+\ell}(q)}{(q^2;q^2)_n} \left(\frac{z}{q^\ell}\right)^{2n} - \frac{(-1)^\ell q^{-\binom{\ell}{2}}}{(q^2, q^3; q^5)_\infty} \sum_{n=0}^{\infty} \frac{b_{2n+\ell}(q)}{(q^2;q^2)_n} \left(\frac{z}{q^\ell}\right)^{2n}.$$

This proves the following theorem.

Theorem 2.1. *If $\ell \in \mathbb{N}_0$, $z \notin \{1, q^{-1}, q^{-2}, \cdots\}$ then*

$$(z;q)_\infty \, {}_1\phi_1\left(-q^{\ell+1}/z; z; q, -z\right)$$

$$= \frac{(-1)^\ell q^{-\binom{\ell}{2}}}{(q, q^4; q^5)_\infty} \sum_{n=0}^{\infty} \frac{a_{2n+\ell}(q)}{(q^2;q^2)_n} \left(\frac{z}{q^\ell}\right)^{2n} - \frac{(-1)^\ell q^{-\binom{\ell}{2}}}{(q^2, q^3; q^5)_\infty} \sum_{n=0}^{\infty} \frac{b_{2n+\ell}(q)}{(q^2;q^2)_n} \left(\frac{z}{q^\ell}\right)^{2n}. \tag{2.6}$$

Let $z = -q^{\ell+1}$ in (1.11) to get

$$\left(-q^{\ell+1};q\right)_\infty = (-1)^\ell q^{-\binom{\ell}{2}} \sum_{n=0}^{\infty} \left[\frac{a_{2n+\ell}(q) q^{2n}}{(q, q^4; q^5)_\infty (q^2;q^2)_n} - \frac{b_{2n+\ell}(q) q^{2n}}{(q^2, q^3; q^5)_\infty (q^2;q^2)_n}\right]. \tag{2.7}$$

From (1.22) to obtain

$$A_q(-b) = \frac{1}{(q, q^4; q^5)_\infty} \sum_{k=0}^{\infty} \frac{(b;q)_k \, q^k a_k(q)}{(q;q)_k} - \frac{1}{(q^2, q^3; q^5)_\infty} \sum_{k=0}^{\infty} \frac{(b;q)_k \, q^k b_k(q)}{(q;q)_k}. \tag{2.8}$$

The special case $b = q^m$ of (2.8) is the interesting identity

$$(-1)^m q^{\binom{m}{2}} \left[\frac{a_m(q)}{(q^1, q^4, q^5; q^5)_\infty} - \frac{b_m(q)}{(q^2, q^3, q^5; q^5}{}_\infty\right]$$

$$= \frac{1}{(q^1, q^4, q^5; q^5)_\infty} \sum_{k=0}^{\infty} \frac{(q^m;q)_k}{(q;q)_k} q^k a_k(q) - \frac{1}{(q^2, q^3, q^5; q^5)_\infty} \sum_{k=0}^{\infty} \frac{(q^m;q)_k}{(q;q)_k} q^k b_k(q). \tag{2.9}$$

The further specialization, $m = 1$, is also interesting

$$1 + \sum_{k=0}^{\infty} b_k(q) \, q^k = \frac{\left(q^2, q^3; q^5\right)_\infty}{(q, q^4; q^5)_\infty} \sum_{k=0}^{\infty} a_k(q) \, q^k. \tag{2.10}$$

The identities (2.9) and (2.10) should have very interesting partition theoretic interpretations.

Theorem 2.2. *For any $m, n \in \mathbb{N}$ we have*

$$(-1)^m q^{-\binom{m}{2}} \left[\frac{a_m(q)}{(q^1, q^4, q^5; q^5)_\infty} - \frac{b_m(q)}{(q^2, q^3, q^5; q^5}{}_\infty\right]$$

$$= (-1)^{m+n} \sum_{k=0}^{\infty} \frac{q^{k(k+m)}}{(q;q)_k} q^{-\binom{m+n+2k}{2}} \left[\frac{a_{m+n+2k}(q)}{(q, q^4, q^5; q^5)_\infty} - \frac{b_{m+n+2k}(q)}{(q^2, q^3, q^5; q^5}{}_\infty\right]. \tag{2.11}$$

Proof. In [23] we proved that

$$A_q(z) = \sum_{k=0}^{\infty} \frac{q^{k^2}}{(q;q)_k} (-z)^k (w;q)_k \, A_q\left(wzq^{2k}\right). \tag{2.12}$$

The theorem is the special case $z = -q^m, w = q^n$ in (2.12). □

It must be noted that the series on the right-hand side of (2.11) converges because (1.2) shows that

$$\lim_{m\to\infty}\left[\frac{(-1)^m q^{-\binom{m}{2}}a_m(q)}{(q,q^4;q^5)_\infty}-\frac{(-1)^m q^{-\binom{m}{2}}b_m(q)}{(q^2,q^3;q^5)_\infty}\right]=1.$$

Using (1.20) and (2.1) we find that for $m=0,1,\cdots$, we have

$$\begin{aligned}(-1)^m q^{-\binom{m}{2}}&\left[\frac{a_m(q)}{(q,q^4;q^5)_\infty}-\frac{b_m(q)}{(q^2,q^3;q^5)_\infty}\right]\\ &=(-q^{m+2};q^2)_\infty\sum_{n=0}^{\infty}\frac{q^{n^2+mn}}{(q^2,-q^{m+2};q^2)_n}\end{aligned}\tag{2.13}$$

Here again the cases $m=0$ and $m=1$ are

$$\frac{1}{(q,q^4;q^5)_\infty}=(-q^2;q^2)_\infty\sum_{n=0}^{\infty}\frac{q^{n^2}}{(q^2,-q^2;q^2)_k},\tag{2.14}$$

$$\frac{1}{(q^2,q^3;q^5)_\infty}=(-q^3;q^2)_\infty\sum_{n=0}^{\infty}\frac{q^{n^2+n}}{(q^2,-q^3;q^2)_n}.\tag{2.15}$$

Neither (2.14) nor (2.15) seem to be on the Slater list [29]

3 Bilateral Sums

In this section we derive bilateral sum identities.

Theorem 3.1. *Let $r\geq s$, $\alpha_j\in\mathbb{C}$, $1\leq j\leq r$, $\beta_k\in\mathbb{C}$, $1\leq k\leq s$, $|\beta_1/\alpha_1|<|w|<1$ and $|z|<|w|$ if $r=s+1$. Then*

$$\begin{aligned}\sum_{m=-\infty}^{\infty}&\frac{(\alpha_1;q)_m}{(\beta_1;q)_m}{}_r\phi_s\left(\begin{matrix}\alpha_1q^m,\ldots,\alpha_r\\ \beta_1q^m,\ldots,\beta_s\end{matrix}\,\middle|\,q,z\right)w^m\\ &=\frac{(q/(\alpha_1w),\alpha_1w,q,\beta_1/\alpha_1;q)_\infty}{(\beta_1,q/\alpha_1,w,\beta_1/(\alpha_1w);q)_\infty}\,{}_{r-1}\phi_{s-1}\left(\begin{matrix}\alpha_2,\ldots,\alpha_r\\ \beta_2,\ldots,\beta_s\end{matrix}\,\middle|\,q,\frac{z}{w}\right)\end{aligned}\tag{3.1}$$

Proof. The left-hand side is

$$\begin{aligned}&=\sum_{m=-\infty}^{\infty}w^m\sum_{n=0}^{\infty}\frac{(\alpha_1;q)_{n+m}}{(\beta_1;q)_{n+m}}\frac{(\alpha_2,\ldots,\alpha_r;q)_n\,z^n}{(q,\beta_2,\ldots,\beta_s;q)_n}\left(-q^{(n-1)/2}\right)^{n(s+1-r)}\\
&=\sum_{n=0}^{\infty}\frac{(\alpha_1,\ldots,\alpha_r;q)_n\,z^n}{(q,\beta_1,\ldots,\beta_s;q)_n}\left(-q^{(n-1)/2}\right)^{n(s+1-r)}\sum_{m=-\infty}^{\infty}\frac{(\alpha_1q^n;q)_m}{(\beta_1q^n;q)_m}w^m\\
&=\sum_{n=0}^{\infty}\frac{(\alpha_1,\ldots,\alpha_r;q)_n\,z^n}{(q,\beta_1,\ldots,\beta_s;q)_n}\left(-q^{(n-1)/2}\right)^{n(s+1-r)}\frac{\left(\beta_1/\alpha_1,q,q^{1-n}(\alpha_1w),\alpha_1wq^n;q\right)_\infty}{(\beta_1q^n,\beta_1/(\alpha_1w),q^{1-n}/\alpha_1,w;q)_\infty}\\
&=\frac{(\beta_1/\alpha_1,\alpha_1w,q;q)_\infty}{(\beta_1,\beta_1/(\alpha_1w),w;q)_\infty}\sum_{n=0}^{\infty}\frac{(\alpha_1,\ldots,\alpha_r;q)_n\,z^n}{(q,\alpha_1w,\beta_2,\ldots,\beta_s;q)_n}\left(-q^{(n-1)/2}\right)^{n(s+1-r)}\frac{\left(q^{1-n}/(\alpha_1w);q\right)_\infty}{(q^{1-n}/\alpha_1;q)_\infty}\\
&=\frac{(q/(\alpha_1w),\alpha_1w,q,\beta_1/\alpha_1;q)_\infty}{(\beta_1,q/\alpha_1,w,\beta_1/(\alpha_1w);q)_\infty}\sum_{n=0}^{\infty}\frac{(\alpha_2,\ldots,\alpha_r;q)_n\,(z/w)^n}{(q,\beta_2,\ldots,\beta_s;q)_n}\left(-q^{(n-1)/2}\right)^{n(s+1-r)}.\end{aligned}$$

and the theorem follows. □

Corollary 3.2. *For $|\beta_1/\alpha_1| < |\alpha_2\alpha_3 z/\beta_2| < 1$ and $|\alpha_2\alpha_3| < |\beta_2|$ we have*

$$\begin{aligned}(3.2)\qquad &\sum_{m=-\infty}^{\infty} \frac{(\alpha_1;q)_m}{(\beta_1;q)_m} {}_3\phi_2\left(\begin{array}{c}\alpha_1 q^m, \alpha_2, \alpha_3 \\ \beta_1 q^m, \beta_2\end{array}\middle| q, z\right)\left(\frac{\alpha_2\alpha_3 z}{\beta_2}\right)^m \\ &= \frac{(q\beta_2/(\alpha_1\alpha_2\alpha_3 z), \alpha_1\alpha_2\alpha_3 z/\beta_2, q, \beta_1/\alpha_1, \beta_2/\alpha_2, \beta_2/\alpha_3); q)_\infty}{(\beta_1, \beta_2, \beta_2/(\alpha_2\alpha_3), q/\alpha_1, \alpha_2\alpha_3 z/\beta_2, \beta_1\beta_2/(\alpha_1\alpha_2\alpha_3 z); q)_\infty}.\end{aligned}$$

Proof. Apply Theorem 3.1 with $w = z\beta_2/\alpha_2\alpha_3$ and use the q-Gauss sum. □

Corollary 3.3. *(i) For $|\beta_1/\alpha_1| < |w| < 1$ and $|z| < |w|$ we have*

$$\begin{aligned}(3.3)\qquad &\sum_{m=-\infty}^{\infty} \frac{(\alpha_1;q)_m}{(\beta_1;q)_m} {}_2\phi_1\left(\begin{array}{c}\alpha_1 q^m, \alpha_2 \\ \beta_1 q^m\end{array}\middle| q, z\right) w^m \\ &= \frac{(q/(\alpha_1 w), \alpha_1 w, q, \beta_1/\alpha_1, a_2 z/w; q)_\infty}{(\beta_1, q/\alpha_1, w, \beta_1/(\alpha_1 w), z/w; q)_\infty}.\end{aligned}$$

(ii) For $|\beta_1/\alpha_1| < |\alpha_2 z/\beta_2| < 1$ we have For $|\beta_1/\alpha_1| < |\alpha_2 z/\beta_2| < 1$ we have

$$\begin{aligned}(3.4)\qquad &\sum_{m=-\infty}^{\infty} \frac{(\alpha_1;q)_m}{(\beta_1;q)_m} {}_2\phi_2\left(\begin{array}{c}\alpha_1 q^m, \alpha_2 \\ \beta_1 q^m, \beta_2\end{array}\middle| q, z\right)\left(\frac{\alpha_2 z}{\beta_2}\right)^m \\ &= \frac{(q/(\alpha_1 w), \alpha_1 w, q, \beta_1/\alpha_1, \beta_2/\alpha_2; q)_\infty}{(\beta_1, \beta_2, q/\alpha_1, w, \beta_1/(\alpha_1 w); q)_\infty}.\end{aligned}$$

Proof. The results follow from Theorem 3.1 and the q-binomial theorem and a confluent limit of the q-Gauss theorem. □

It is important we use the following definition of q-shifted factorial,

$$(3.5)\qquad (a;q)_\infty = \prod_{k=0}^{\infty}\left(1 - aq^k\right), (a;q)_n = \frac{(a;q)_\infty}{(aq^n;q)_\infty},$$

where $a, q, n \in \mathbb{C}$, $|q| < 1$, and $aq^n \neq q^{-k},\ k \in \mathbb{N}_0$, then it is clear that

$$(3.6)\qquad (a;q)_{n+m} = (a;q)_n (aq^n;q)_m, \quad a, n, m \in \mathbb{C}.$$

In the rest of this section we derive identities involving $I_\nu^{(2)}(z;q)$. The next theorem uses Ramanujan's ${}_1\psi_1$ sum [12, (II.28)]

$$(3.7)\qquad \sum_{-\infty}^{\infty} \frac{(a;q)_n}{(b;q)_n} z^n = \frac{(q, b/a, az, q/az; q)_\infty}{(b, q/a, z, b/az; q)_\infty}, \quad |b/a| < |z| < 1.$$

Theorem 3.4. *If $|q^{\nu+1}/a| < |w| < 1$, then*

$$\begin{aligned}(3.8)\qquad &\sum_{m=-\infty}^{\infty} (a;q)_m \left(\frac{w}{z}\right)^m I_{m+\nu}^{(2)}(2z;q) \\ &= \frac{z^\nu \left(q^{\nu+1}/a, aw, q/aw; q\right)_\infty}{(q/a, q^{\nu+1}/aw, w; q)_\infty} \sum_{n=0}^{\infty}\left(-\frac{z^2 q^{\nu+1}}{aw}\right)^n q^{\binom{n}{2}} \frac{(w;q)_n}{(q, q^{\nu+1}/a; q)_n} \\ &= \frac{z^\nu \left(q^{\nu+1}/a, aw, q/aw; q\right)_\infty}{(q/a, q^{\nu+1}/aw, w; q)_\infty} {}_1\phi_1(w; q^{\nu+1}/a; q, z^2 q^{\nu+1}/aw).\end{aligned}$$

Proof. The definition (1.16) implies that

$$I^{(2)}_{m+\nu}(z;q)=\frac{(z/2)^{\nu}}{(q;q)_{\infty}}\sum_{n=0}^{\infty}\frac{q^{n^2+n\nu}}{(q;q)_n}\left(\frac{z^2}{4}\right)^n\left(q^{n+\nu+1};q\right)_{\infty}\frac{\left(zq^n/2\right)^m}{\left(q^{n+\nu+1};q\right)_m}.$$

Assuming $|q^{\nu+1}/a|<|wz/2|<1$ we use the above expansion and establish the generating function

$$\begin{aligned}
&\sum_{m=-\infty}^{\infty}(a;q)_m w^m I^{(2)}_{m+\nu}(z;q)=\frac{(z/2)^{\nu}}{(q;q)_{\infty}}\sum_{n=0}^{\infty}\frac{q^{n^2+n\nu}}{(q;q)_n}\left(\frac{z^2}{4}\right)^n\left(q^{n+\nu+1};q\right)_{\infty}\\
&\times\sum_{m=-\infty}^{\infty}\frac{(a;q)_m\left(wzq^n/2\right)^m}{\left(q^{n+\nu+1};q\right)_m}=\frac{(z/2)^{\nu}}{(q;q)_{\infty}}\sum_{n=0}^{\infty}\frac{q^{n^2+n\nu}}{(q;q)_n}\left(\frac{z^2}{4}\right)^n\left(q^{n+\nu+1};q\right)_{\infty}\\
&\times\frac{\left(q,q^{n+\nu+1}/a,awzq^n/2,2q/(awzq^n);q\right)_{\infty}}{\left(q^{n+\nu+1},q/a,wzq^n/2,2q^{\nu+1}/(awz);q\right)_{\infty}}\\
&=\frac{(z/2)^{\nu}\left(q^{\nu+1}/a,awz/2,2q/(awz);q\right)_{\infty}}{\left(q/a,2q^{\nu+1}/(awz),wz/2;q\right)_{\infty}}\sum_{n=0}^{\infty}\left(-\frac{zq^{\nu+1}}{2aw}\right)^n q^{\binom{n}{2}}\frac{(wz/2;q)_n}{\left(q,q^{\nu+1}/a;q\right)_n},
\end{aligned}$$

where the Ramanujan ${}_1\psi_1$ was used in the second to last step. This establishes our theorem. □

4 Series Involving A_q and S_n

In this section we prove the identities contained in Theorems 4.1 and 4.2 and consider some of their corollaries. The proofs uses the identities

$$S_n\left(ab;q\right) = b^n\sum_{k=0}^{n}\frac{\left(b^{-1};q\right)_k(-1)^k q^{-nk}q^{\binom{k+1}{2}}}{(q;q)_k}S_{n-k}\left(aq^k;q\right), \tag{4.1}$$

$$S_{2n+1}\left(q^{-2n-1};q\right) = 0,\quad S_{2n}\left(q^{-2n};q\right)=\frac{(-1)^n q^{-n^2}}{\left(q^2;q^2\right)_n}. \tag{4.2}$$

$$S_n\left(-q^{-n+1/2};q\right) = \frac{q^{-\binom{n^2-n}{4}/4}}{\left(q^{1/2};q^{1/2}\right)_n},\quad S_n\left(-q^{-n-1/2};q\right)=\frac{q^{-\binom{n^2+n}{4}/4}}{\left(q^{1/2};q^{1/2}\right)_n}. \tag{4.3}$$

These identities were proved in Section 6 of our paper [22].

Theorem 4.1. *For $n=0,1,\dots$ we have*

$$\frac{q^{\binom{n}{2}}S_n\left(bq^{-n};q\right)}{(-b)^n} = \sum_{j=0}^{\lfloor n/2\rfloor}\frac{q^{j^2-j}(-1)^j}{\left(q^2;q^2\right)_j}\frac{\left(b^{-1};q\right)_{n-2j}}{(q;q)_{n-2j}}, \tag{4.4}$$

$$\frac{S_n\left(-q^{-n+1/2}b;q\right)q^{\binom{n}{2}}}{(-b)^n} = \sum_{k=0}^{n}\frac{q^{\binom{k^2-k}{4}/4}}{\left(q^{1/2};q^{1/2}\right)_k}\frac{\left(b^{-1};q\right)_{n-k}}{(q;q)_{n-k}}(-1)^k, \tag{4.5}$$

$$\frac{S_n\left(-q^{-n-1/2}b;q\right)q^{\binom{n}{2}}}{(-b)^n} = \sum_{k=0}^{n}\frac{q^{\binom{k^2-3k}{4}/4}}{\left(q^{1/2};q^{1/2}\right)_k}\frac{\left(b^{-1};q\right)_{n-k}}{(q;q)_{n-k}}(-1)^k, \tag{4.6}$$

Proof. The proof of (4.4) follows from (4.1) and (4.2). Formulas (4.5) and (4.6) follow from (4.1) and both parts of (4.3). □

The proof of the next theorem makes use of

$$I^{(2)}_{\nu}\left(2q^{-n/2};q\right)=\frac{q^{\nu n/2}S_n\left(-q^{-\nu-n};q\right)}{\left(q^{n+1};q\right)_{\infty}}=\frac{q^{-\nu n/2}S_n\left(-q^{\nu-n};q\right)}{\left(q^{n+1};q\right)_{\infty}}, \tag{4.7}$$

from our work [22].

Theorem 4.2. *The Ramanujan function A_q satisfies the following summation theorems*

$$\sum_{k=0}^{\infty} \frac{q^{\binom{k+1}{2}} A_q\left(aq^{k-2n}\right)}{(q;q)_k} = (-q;q)_\infty (aq^{1-2n};q^2)_\infty, \tag{4.8}$$

$$\sum_{k=0}^{\infty} \frac{q^{\binom{k}{2}} \left(-q^{3/2}\right)^k A_q\left(-q^{k-n+1/2}\right)}{(q;q)_k} = \frac{(q;q)_n}{q^{(n^2-n)/4}\left(q^{1/2};q^{1/2}\right)_n}, \tag{4.9}$$

$$\sum_{k=0}^{\infty} \frac{q^{k^2/2} (-1)^k A_q\left(-q^{k-n-1/2}\right)}{(q;q)_k} = \frac{(-1)^n (q;q)_n}{q^{(n^2+n)4}\left(q^{1/2};q^{1/2}\right)_n}. \tag{4.10}$$

Moreover we have the expansions

$$A_q(w) = (wq;q)_\infty \sum_{n=0}^{\infty} \frac{q^{3n^2}\left(-w^2\right)^n}{\left(q^2,wq,wq^2;q^2\right)_n}. \tag{4.11}$$

$$A_q(w) = \left(-wq^{1/2};q\right)_\infty \sum_{n=0}^{\infty} \frac{q^{n(3n-1)/4}(-w)^n}{\left(q^{1/2};q^{1/2}\right)_n \left(-wq^{1/2};q\right)_n}, \tag{4.12}$$

$$A_q(w) = \left(-wq^{3/2};q\right)_\infty \sum_{n=0}^{\infty} \frac{q^{n(3n+1)/4}(-w)^n}{\left(q^{1/2};q^{1/2}\right)_n \left(-wq^{3/2};q\right)_n}. \tag{4.13}$$

The special cases $a=1$ and $a=q$ of (4.8) are worth recording. For $n>0$ they are

$$\sum_{k=0}^{\infty} \frac{q^{\binom{k+1}{2}} A_q\left(q^{k-2n-1}\right)}{(q;q)_k} = 0, \tag{4.14}$$

$$\sum_{k=0}^{\infty} \frac{q^{\binom{k+1}{2}} A_q\left(q^{k-2n}\right)}{(q;q)_k} = (-1)^n q^{-n^2}\left(q;q^2\right)_n, \tag{4.15}$$

Proof of Theorem of 4.2. The left-hand side of (4.8) is

$$\sum_{k,m=0}^{\infty} \frac{(-a)^m q^{m^2+\binom{k+1}{2}}}{(q;q)_k (q;q)_m} q^{km-2nm} = \sum_{m=0}^{\infty} \frac{(-a)^m q^{m^2-2mn}}{(q;q)_m} (-q^{m+1};q)_\infty$$
$$= (-q;q)_\infty \sum_{m=0}^{\infty} \frac{(-a)^m q^{m^2-2mn}}{(q^2;q^2)_m},$$

and we have proved (4.8). Let us first rewrite (1.18) in the equivalent form

$$I_\nu^{(2)}(2z;q) = \frac{z^\nu}{(q;q)_\infty} \sum_{k=0}^{\infty} \frac{\left(-q^\nu\right)^k}{(q;q)_k} q^{\binom{k+1}{2}} A_q\left(-q^{\nu+k}z^2\right). \tag{4.16}$$

Formulas (4.9) and (4.10) are the special cases $\nu = 1/2, -1/2$, respectively, combined with (4.7). We now come to (4.11). Write $(wq;q)_\infty/(qw;q)_{2n}$ as $(wq^{2n+1};q)_\infty$ then expand it into powers of w to see that the right-hand side of (4.11) is

$$\sum_{k,n=0}^{\infty} \frac{(-1)^{n+k} w^{2n+k}}{(q;q)_k (q^2;q^2)_n} q^{3n^2+2nk+\binom{k+1}{2}} = \sum_{m=0}^{\infty} (-w)^m q^{\binom{m+1}{2}} \sum_{n=0}^{\lfloor n/2 \rfloor} \frac{(-1)^n q^{n(n-1)}}{(q^2;q^2)_n (q;q)_{m-2n}}.$$

Denote the n-sum by c_m. It is easy to see that

$$\sum_{m=0}^{\infty} c_m t^m = \frac{(t^2;q^2)_\infty}{(t;q)_\infty} = (-t;q)_\infty.$$

Thus $c_m = q^{\binom{m}{2}}/(q;q)_m$. This establishes (4.11). The proof of (4.12) is similar. Its right-hand side is

$$\sum_{n=0}^{\infty} \frac{q^{n(3n-1)/4}\,(-w)^n}{\left(q^{1/2};q^{1/2}\right)_n}(-wq^{n+1/2};q)_\infty = \sum_{n,k=0}^{\infty} \frac{q^{n(3n-1)/4}w^{n+k}}{\left(q^{1/2};q^{1/2}\right)_n}\frac{(-1)^n}{(q;q)_k}q^{nk+k^2/2}$$
$$= \sum_{m=0}^{\infty} w^m q^{m^2/2} \sum_{n+k=m} \frac{(-1)^n q^{n(n-1)/4}}{(q^{1/2};q^{1/2})_n}\,\frac{1}{(q;q)_k}.$$

Denoting the inner sum by d_m we find that

$$\sum_{m=0}^{\infty} d_m t^m = \frac{(t;q^{1/2})_\infty}{(t;q)_\infty} = (tq^{1/2};q)_\infty,$$

and we conclude that $d_m = q^{m^2/2}(-1)^m/(q;q)_m$ and the proof of (4.12) is complete. The proof of (4.13) is parallel to the proof of (4.12) and will be omitted. □

The cases $w = -q^m$ of (4.11)-(4.13) are interesting. They are

$$\frac{(-1)^m q^{-\binom{m}{2}} a_m(q)}{(q,q^4;q^5)_\infty} - \frac{(-1)^m q^{-\binom{m}{2}} b_m(q)}{(q^2,q^3;q^5)_\infty}$$
$$= (-q^{m+1};q)_\infty \sum_{n=0}^{\infty} \frac{(-1)^n q^{3n^2+2mn}}{(q^2;q^2)_n(-q^{m+1};q)_n}, \tag{4.17}$$

$$\frac{(-1)^m q^{-m(m-1)} a_m(q^2)}{(q^2,q^8;q^{10})_\infty} - \frac{(-1)^m q^{-m(m-1)} b_m(q^2)}{(q^4,q^6;q^{10})_\infty}$$
$$= (q^{2m+1};q)_\infty \sum_{n=0}^{\infty} \frac{q^{2mn+n(3n+1)/2}}{(q;q)_n(q^{2m+1};q^2)}_\infty$$
$$= (q^{2m+3};q)_\infty \sum_{n=0}^{\infty} \frac{q^{2mn+n(3n+1)/2}}{(q;q)_n(q^{2m+3};q^2)}_\infty. \tag{4.18}$$

The case $m = 0$ of (4.18) is on the Slater list, see (44)–(45) in [29].

The structure of the identities (4.12) and (4.13) suggests that we consider the series

$$(-wq^{\alpha+1/2};q)_\infty \sum_{n=0}^{\infty} \frac{q^{n(3n+4c)/4}\,(-w)^n}{\left(q^{1/2};q^{1/2}\right)_n\left(-wq^{\alpha+1/2};q\right)_n}$$
$$= \sum_{n,k=0}^{\infty} \frac{(-1)^n q^{n(3n+4c)/4} w^{n+k}}{\left(q^{1/2};q^{1/2}\right)_n (q;q)_k} q^{k(\alpha+n)+k^2/2}$$
$$= \sum_{m=0}^{\infty} w^m q^{\alpha m+m^2/2} \sum_{n=0}^{m} \frac{(-1)^n q^{n(c-\alpha)+n^2/4}}{(q^{1/2};q^{1/2})_n (q;q)_{m-n}}$$

Denote the n sum by c_m. Thus

$$\sum_{m=0}^{\infty} c_m t^m = \frac{(tq^{c-\alpha+1/4};q^{1/2})_\infty}{(t;q)_\infty}.$$

At this stage we take $c = \alpha - 1/4$, or $c = \alpha - 3/4$ and find that $c_m = (-1)^m q^{m^2/2}/(q;q)_m$, or $c_m = (-1)^m q^{m(m-1)/2}/(q;q)_m$, respectively. This essentially leads to (4.12) and (4.13). On the hand hand we have other choices for c.

We let $c=\alpha-1/4-j/2$. In this case it turned out that there is no loss of generality in assuming $\alpha=0$. We also assume $j>0$, since $j=0$ is already covered. Thus

$$\begin{aligned}&(-wq^{1/2};q)_\infty\sum_{n=0}^{\infty}\frac{q^{n(3n-2j-1)/4}\left(-w\right)^n}{\left(q^{1/2};q^{1/2}\right)_n\left(-wq^{1/2};q\right)_n}\\&=\sum_{m=0}^{\infty}w^mq^{m^2/2}\sum_{n=0}^{m}\frac{(-1)^nq^{-jn/2+n(n-1)/4}}{(q;q)_k(q^{1/2};q^{1/2})_n}.\end{aligned}$$

Again we denote the n-sum by c_m and find that

$$\sum_{m=0}^{\infty}c_mt^m=\frac{(tq^{-j/2};q^{1/2})_\infty}{(t;q)_\infty}=(tq^{-j/2};q^{1/2})_{j-1}(tq^{-1/2};q)_\infty.$$

Therefore

$$c_m=(-1)^m\sum_{s+k=m}\begin{bmatrix}j-1\\s\end{bmatrix}_{q^{1/2}}q^{s(s-1)/4}q^{-js/2}\frac{q^{k\binom{k-2}{}/2}}{(q;q)_k}.$$

This proves that

$$\begin{aligned}&(-wq^{1/2};q)_\infty\sum_{n=0}^{\infty}\frac{q^{n(3n-2j-1)/4}\left(-w\right)^n}{\left(q^{1/2};q^{1/2}\right)_n\left(-wq^{1/2};q\right)_n}\\&=\sum_{s=0}^{j}\begin{bmatrix}j-1\\s\end{bmatrix}_{q^{1/2}}q^{s(3s-2j-1)/4}w^sA_q(wq^{s-1}).\end{aligned}\tag{4.19}$$

It is clear that (4.13) is the special case $j=1$, with $w\to wq$.

Lemma 4.3. *The Stieltjes-Wigert polynomials satisfy the generating function*

$$(z,zt;q)_\infty=\sum_{k=0}^{\infty}\left(-z\right)^kq^{\binom{k}{2}}S_k\left(-tq^{-k};q\right).\tag{4.20}$$

For $n=0,1,\dots$ we have following evaluations

$$S_{2n+1}\left(q^{-2n-1};q\right)=0,\quad S_{2n}\left(q^{-2n};q\right)=\frac{\left(-1\right)^nq^{n-n^2}}{\left(q^2;q^2\right)_n}.\tag{4.21}$$

and

$$S_n\left(-q^{-n+1/2};q\right)=\frac{q^{-\left(n^2-n\right)/4}}{\left(q^{1/2};q^{1/2}\right)_n},\ S_n\left(-q^{-n-1/2};q\right)=\frac{q^{-\left(n^2+n\right)/4}}{\left(q^{1/2};q^{1/2}\right)_n}.\tag{4.22}$$

Proof. From (1.7) and (1.8) we get,

$$\begin{aligned}&\left(-zq^{1/2},zt;q\right)_\infty=\frac{1}{\sqrt{2\pi\log q^{-1}}}\int_{-\infty}^{\infty}\frac{(zt;q)_\infty\exp\left(\frac{x^2}{\log q^2}\right)}{(ze^{ix};q)_\infty}dx\\&=\sum_{k=0}^{\infty}z^k\int_{-\infty}^{\infty}\frac{\left(te^{-ix};q\right)_k\exp\left(\frac{x^2}{\log q^2}+ikx\right)}{\sqrt{2\pi\log q^{-1}}\left(q;q\right)_k}dx=\sum_{k=0}^{\infty}z^k\int_{-\infty}^{\infty}\frac{\left(te^{ix};q\right)_k\exp\left(\frac{x^2}{\log q^2}-ikx\right)}{\sqrt{2\pi\log q^{-1}}\left(q;q\right)_k}dx\\&=\sum_{k=0}^{\infty}z^kq^{k^2/2}S_k\left(tq^{-k-1/2};q\right).\end{aligned}$$

Let $t = q^{1/2}$ in (4.20) to get (4.21) by matching the corresponding coefficients of z^n of

$$\left(z^2q;q^2\right)_\infty = \sum_{n=0}^{\infty} \frac{q^{n^2}(-1)^n z^{2n}}{(q^2;q^2)_n} = \sum_{k=0}^{\infty} (-z)^k \, q^{\binom{k}{2}} S_k\left(q^{-k};q\right).$$

Let $t = -1$ and $t = -q$ in (4.20) to obtain

$$\left(-zq^{1/2},-z;q\right)_\infty = \left(-z;q^{1/2}\right)_\infty = \sum_{n=0}^{\infty} \frac{q^{n(n-1)/4}z^n}{\left(q^{1/2};q^{1/2}\right)_n} = \sum_{k=0}^{\infty} z^k q^{k^2/2} S_k\left(-q^{-k-1/2};q\right),$$

$$\left(-zq^{1/2},-zq;q\right)_\infty = \left(-zq^{1/2};q^{1/2}\right)_\infty = \sum_{n=0}^{\infty} \frac{q^{n(n+1)/4}z^n}{\left(q^{1/2};q^{1/2}\right)_n} = \sum_{k=0}^{\infty} z^k q^{k^2/2} S_k\left(-q^{-k+1/2};q\right)$$

respectively, then evaluations in (4.22) are obtained by matching corresponding coefficients of z^n. □

Theorem 4.4. *For $|c/(ab)| < 1$ we have*

$$\sum_{n=0}^{\infty} \frac{(b;q)_n \, q^{n^2/2} S_n\left(aq^{-n-1/2};q\right)}{(c;q)_n} \left(\frac{c}{ab}\right)^n = \frac{(c/b;q)_\infty}{(c;q)_\infty} \sum_{n=0}^{\infty} \frac{q^{n^2/2}}{(q;q)_n} \left(\frac{c}{ab}\right)^n A_q\left(\frac{cq^{n-1/2}}{a}\right). \tag{4.23}$$

In particular, for $|cz| < q^{1/2}$ we have

$$\sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}}}{(q;q)_n} (cz)^n A_q\left(cq^{n-1}\right) = \frac{(c;q)_\infty}{(cz;q)_\infty} \sum_{n=0}^{\infty} \frac{q^{n^2} \prod_{j=0}^{2n-1}\left(z-q^j\right)\left(-c^2\right)^n}{\left(q^2,c,cq;q^2\right)_n} \tag{4.24}$$

and

$$A_q(c) = \sum_{n=0}^{\infty} \frac{q^{3n^2+n}\left(-c^2\right)^n}{\left(q^2,cq,cq^2;q^2\right)_n}. \tag{4.25}$$

Similarly, for $\left|c^2z\right| < q^{-1}$ we have

$$\sum_{n=0}^{\infty} \frac{q^{n^2+n}\left(-c^2z\right)^n}{\left(q^2;q^2\right)_n} A_{q^2}\left(-c^2q^{2n}\right) = \frac{\left(c^2q;q^2\right)_\infty}{\left(c^2zq;q^2\right)_\infty} \sum_{n=0}^{\infty} \frac{q^{\binom{n+1}{2}} \prod_{j=0}^{n-1}\left(z-q^{2j}\right)\left(-c^2\right)^n}{\left(q,cq^{1/2},-cq^{1/2};q\right)_n} \tag{4.26}$$

and

$$A_{q^2}\left(-c^2\right) = \left(c^2q;q^2\right)_\infty \sum_{n=0}^{\infty} \frac{q^{(3n^2-n)/2}c^{2n}}{\left(q,cq^{1/2},-cq^{1/2};q\right)_n}. \tag{4.27}$$

Proof. Let $\alpha = -n$ in (4.20) to get

$$q^{n^2/2} S_n\left(xq^{-n-1/2};q\right) = \frac{1}{\sqrt{\pi \log q^{-2}}} \int_{-\infty}^{\infty} \frac{\left(xe^{iy};q\right)_n}{(q;q)_n} \exp\left(\frac{y^2}{\log q^2} - iny\right) dy.$$

Under the condition $|c/(ab)| < 1$ we first apply the q-Gauss sum (II.8) of [12] then apply the

q-binomial theorem (II.1) of [12] to obtain

$$\sum_{n=0}^{\infty}\frac{(b;q)_n\,q^{n^2/2}S_n\left(aq^{-n-1/2};q\right)}{(c;q)_n}\left(\frac{c}{ab}\right)^n=\int_{-\infty}^{\infty}\sum_{n=0}^{\infty}\frac{\left(b,ae^{iy};q\right)_n}{(q,c;q)_n}\left(\frac{ce^{-iy}}{ab}\right)^n\frac{\exp\left(\frac{y^2}{\log q^2}\right)dy}{\sqrt{\pi\log q^{-2}}}$$
$$=\int_{-\infty}^{\infty}\frac{\left(ce^{-iy}/a,c/b;q\right)_\infty}{\left(c,ce^{-iy}/ab;q\right)_\infty}\frac{\exp\left(\frac{y^2}{\log q^2}\right)dy}{\sqrt{\pi\log q^{-2}}}=\int_{-\infty}^{\infty}\frac{\left(ce^{iy}/a,c/b;q\right)_\infty}{\left(c,ce^{iy}/ab;q\right)_\infty}\frac{\exp\left(\frac{y^2}{\log q^2}\right)dy}{\sqrt{\pi\log q^{-2}}}$$
$$=\frac{(c/b;q)_\infty}{(c;q)_\infty}\sum_{n=0}^{\infty}\frac{1}{(q;q)_n}\left(\frac{c}{ab}\right)^n\int_{-\infty}^{\infty}\left(ce^{iy}/a;q\right)_\infty\frac{\exp\left(\frac{y^2}{\log q^2}+iny\right)dy}{\sqrt{\pi\log q^{-2}}}$$
$$=\frac{(c/b;q)_\infty}{(c;q)_\infty}\sum_{n=0}^{\infty}\frac{q^{n^2/2}}{(q;q)_n}\left(\frac{c}{ab}\right)^n A_q\left(\frac{cq^{n-1/2}}{a}\right),$$

which gives (4.23).

Let $a=q^{1/2}$ in (4.23), under the condition $|c/b|<q^{1/2}$ by applying (4.21) to get

$$\frac{(c/b;q)_\infty}{(c;q)_\infty}\sum_{n=0}^{\infty}\frac{q^{\binom{n}{2}}}{(q;q)_n}\left(\frac{c}{b}\right)^n A_q\left(cq^{n-1}\right)=\sum_{n=0}^{\infty}\frac{(b;q)_n\,q^{\binom{n}{2}}S_n\left(q^{-n};q\right)}{(c;q)_n}\left(\frac{c}{b}\right)^n$$
$$=\sum_{n=0}^{\infty}\frac{(b;q)_{2n}\,q^{\binom{2n}{2}}S_{2n}\left(q^{-2n};q\right)}{(c;q)_{2n}}\left(\frac{c}{b}\right)^{2n}=\sum_{n=0}^{\infty}\frac{(-1)^n\,(b;q)_{2n}\,q^{n^2}}{(c;q)_{2n}\,(q^2;q^2)_n}\left(\frac{c}{b}\right)^{2n}$$
$$=\sum_{n=0}^{\infty}\frac{q^{n^2}\prod_{j=0}^{2n-1}\left(1/b-q^j\right)\left(-c^2\right)^n}{(c;q)_{2n}\,(q^2;q^2)_n}=\sum_{n=0}^{\infty}\frac{q^{n^2}\prod_{j=0}^{2n-1}\left(1/b-q^j\right)\left(-c^2\right)^n}{(q^2,c,cq;q^2)_n},$$

which gives (4.24). Let $b\to\infty$ in the above equation or $z=0$ in (4.24) to obtain (4.25). Let $a=-1$ in (4.23), by (4.22) to obtain

$$\frac{(c/b;q)_\infty}{(c;q)_\infty}\sum_{n=0}^{\infty}\frac{q^{n^2/2}}{(q;q)_n}\left(-\frac{c}{b}\right)^n A_q\left(-cq^{n-1/2}\right)=\sum_{n=0}^{\infty}\frac{(b;q)_n\,q^{n^2/2}S_n\left(-q^{-n-1/2};q\right)}{(c;q)_n}\left(-\frac{c}{b}\right)^n$$
$$=\sum_{n=0}^{\infty}\frac{(b;q)_n\,q^{\left(n^2-n\right)/4}}{(c;q)_n\left(q^{1/2};q^{1/2}\right)_n}\left(-\frac{c}{b}\right)^n=\sum_{n=0}^{\infty}\frac{q^{\binom{n}{2}/2}\prod_{j=0}^{n-1}\left(1/b-q^j\right)(-c)^n}{\left(q^{1/2},c^{1/2},-c^{1/2};q^{1/2}\right)_n},$$

which gives (4.26), taking $b\to\infty$ in the above equation to get (4.27). □

Corollary 4.5. *For $m=0,1,\ldots,$ we have*

$$\sum_{n=0}^{\infty}\frac{q^{3n^2+(2m+1)n}\,(-1)^n}{\left(q^2,-q^{m+1},-cq^{m+2};q^2\right)_n}=\frac{(-1)^m q^{-\binom{m}{2}}a_m(q)}{(q,q^4;q^5)_\infty}-\frac{(-1)^m q^{-\binom{m}{2}}b_m(q)}{(q^2,q^3;q^5)_\infty},\tag{4.28}$$

and

$$\sum_{n=0}^{\infty}\frac{q^{(3n^2-n)/2+2mn}}{\left(q,q^{m+1/2},-q^{m+1/2};q\right)_n}=\frac{(-1)^m q^{-m(m-1)}}{\left(q^{2m+1};q^2\right)_\infty}\left\{\frac{a_m(q^2)}{(q^2,q^8;q^{10})_\infty}-\frac{b_m(q^2)}{(q^4,q^6;q^{10})_\infty}\right\}.\tag{4.29}$$

Proof. Equation (4.28) is obtained by letting $c=-q^m,\ m=0,1,\ldots$ in (4.25), while Equation (4.29)is followed by setting $c=q^m,\ m=0,1,\ldots$ in (4.27). □

References

[1] N. I. Akhiezer, *The Classical Moment Problem and Some Related Questions in Analysis*, English translation, Oliver and Boyd, Edinburgh, 1965

[2] G. E. Andrews, *The Theory of Partitions*, Addison-Wesley, Reading, MA, 1976.

[3] G. E. Andrews, *q-series: Their development and application in analysis, number theory, combinatorics, physics, and computer algebra*, CBMS Regional Conference Series, Number 66, American Mathematical Society, Providence, RI, 1986.

[4] G. E. Andrews, A polynomial identity which implies the Rogers–Ramanujan identities, Scripta Math. **28** (1970), 297–305.

[5] G. E. Andrews, R. A. Askey, and R. Roy, *Special Functions*, Cambridge University Press, Cambridge, 1999.

[6] R. A. Askey, Continuous q-Hermite polynomials when $q > 1$, in "q-Series and Partitions", D. Stanton, ed., IMA Volumes in Mathematics and Its Applications, Springer-Verlag, New York, 1989, pp. 151–158.

[7] L. Carlitz, A note on the Rogers–Ramanujan identities, Math. Nachrichten, **17** (1958), 23–26.

[8] J. S. Christiansen, The moment problem associated with the Stieltjes–Wigert polynomials, J. Math. Anal. Appl. **277** (2003), 218245.

[9] J. S. Christiansen, The moment problem associated with the q-Laguerre polynomials, Constr. Approx. **19** (2003), 122.

[10] A. Erdélyi, W. Magnus, F. Oberhettinger, F. G. Tricomi, *Tables of integral transforms*, vol. I, McGraw-Hill Book Company, , New York, 1954.

[11] K. Garrett, M. E. H. Ismail, and D. Stanton, Variants of the Rogers–Ramanujan identities, Advances in Applied Math. **23** (1999), 274–299.

[12] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, second edition Cambridge University Press, Cambridge, 2004.

[13] M. E. H. Ismail, The zeros of basic Bessel functions, the functions $J_{\nu+ax}(x)$ and the associated orthogonal polynomials, J. Math. Anal. Appl. bf 86 (1982), 1–19.

[14] M. E. H. Ismail, Asymptotics of q orthogonal polynomials and a q-Airy function, IMRN (2005), 1063–1088.

[15] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in one variable*, Cambridge University Press, paperback edition, Cambridge, 2009.

[16] M. E. H. Ismail and D. R. Masson, q-Hermite polynomials, biorthogonal rational functions, and q-beta integrals, Transactions Amer. Math. Soc. **346** (1994), 63–116.

[17] M. E. H. Ismail and M. Rahman, The q-Laguerre polynomials and related moment problems. J. Math. Anal. Appl. **218** (1998), 155174.

[18] M. E. H. Ismail and D. R. Stanton, Tribasic Integrals and identities of Rogers-Ramanujan type, Transactions Amer. Math. Soc. **355** (2003), 4061-4091.

[19] M. E. H. Ismail and C. Zhang, Zeros of entire functions and a problem of Ramanujan, Advances in Math. **209** (2007), 363–380.

[20] M. E. H. Ismail and R. Zhang, Diagonalization of certain integral operators, Advances in Math. **109** (1994), 1–33.

[21] M. E. H. Ismail and R. Zhang, Chaotic and periodic asymptotics for q-orthogonal polynomials, IMRN, Article ID 83274, (2006), 1–33.

[22] M. E. H. Ismail and R. Zhang, q-Bessel Functions and Rogers-Ramanujan Type Identities, Proc. Amer. Math. Soc. (2017), to appear.

[23] M. E. H. Ismail and R. Zhang, Integral and series representations of q-polynomials and functions: part I, to appear.

[24] R. Koekoek and R. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogues, Reports of the Faculty of Technical Mathematics and Informatics no. 98-17, Delft University of Technology, Delft, 1998.

[25] H. Rademacher, *Topics in Analytic Number Theory*, Edited by E. Grosswald, J. Lehner, and M. Newman. Die Grundlehren der mathematischen Wissenschaften, Band 169, Springer-Verlag, New York-Heidelberg, 1973.

[26] S. Ramanujan, The Lost Notebook and Other Unpublished Papers (Introduction by G. E. Andrews), Narosa, New Delhi, 1988.

[27] S. Ramanujan, *Collected Papers of Sirinivasa Ramanujan*, G. H. Hardy, P. V. Seshu Aiyar, and B. M. WIlson; editors, AMS Chelsea Publications, American Mathematical Society, Providence, Rhode Island, 2000.

[28] I. Schur, "Ein Beitrag zur Additiven Zahlentheorie und zur Theorie der Kettenbrüche, Sitzungsberichte der Preussischen Akademie der Wissenschaften, physikalish??mathema- tische Klasse, pp. 302–??321, 1917; reprinted in I. Schur, Gesmmelte Abhandungen,?? Vol. 2, pp. 117–136, Springer-Verlag, Berlin, 1973.

[29] L. J. Slater, Further identies of the Rogers-Ramanujan type. Proc. London Math. Soc. (2) **54** (1952), 147–167.

[30] G. Szegő, *Orthogonal Polynomials*, fourth edition, American Mathematical Society, Providence, 1975.

[31] E. C. Titchmarsh, *Introduction to the Theory of Fourier Integrals*, second edition, Oxford University Press, Oxford, 1948.

[32] E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, fourth edition, Cambridge University Press, Cambridge, 1927.

[33] R. Zhang, Scaled asymptotics for q-series, Quarterly J. Math. Oxford **59** (2008), 389–407.

M. E. H. I, University of Central Florida, Orlando, Florida 32816
and King Saud University, Riyadh, Saudi Arabia
email: ismail@math.ucf.edu

R. Z., College of Science, Northwest A&F University, Yangling, Shaanxi 712100, P. R. China PRC
email: ruimingzhang@yahoo.com